arXiv:1407.1232v1 [cs.IT] 19 Jun 2014

On Quantum Codes Obtained From Cyclic Codes Over $F_2 + vF_2 + v^2F_2$

Abdullah Dertli[a], Yasemin Cengellenmis[b], Senol Eren[a]
(a) Ondokuz Mayıs University, Faculty of Arts and Sciences,
Mathematics Department, Samsun, Turkey
abdullah.dertli@gmail.com, seren@omu.edu.tr
(b) Trakya University, Faculty of Arts and Sciences,
Mathematics Department, Edirne, Turkey
ycengellenmis@gmail.com

July 7, 2014

Abstract

A new Gray map which is both an isometry and a weight preserving map from $R = F_2 + vF_2 + v^2F_2$ to F_2^3 is defined. A construction for quantum error correcting codes from cyclic codes over finite ring $R = F_2 + vF_2 + v^2F_2$, $v^3 = v$ is given.The parameters of quantum codes which are obtained from cyclic codes over R are determined.

1 Introduction

Quantum computer has extraordinary power and it can certain hard problems much more quickly than any classical computer. It is a computation device that makes direct use of quantum mechanical phenomena to perform on data.

Quantum computer inevitably interacts with its surrounding, resulting in decoherence and hence in the decay of quantum information stored in the device.

Quantum error correction is used in quantum computing to protect quantum information from errors due to decoherence and other quantum noise. Effective techniques for quantum error correction were first developed by Shor and independently by Steane. They discovered quantum error correcting codes.

Although the theory quantum error correcting codes has striking differences from the theory classical error correcting codes, Calderbank et al. gave a way to construct quantum error correcting codes from classical error correcting codes.

Many good quantum codes have been constructed by using classical cyclic codes over F_q with self orthogonal (or dual containing) properties.

Some authors constructed quantum codes by using linear codes over finite rings. For example, in [4], J.Qian et al. gave a new method to obtain self-orthogonal codes over F_2. They gave a construction for quantum error correcting

codes starting from cyclic codes over finite ring, $F_2 + uF_2$, $u^2 = 0$. X.Kai, S.Zhu gave construction for quantum codes from linear and cyclic codes over $F_4 + uF_4$, $u^2 = 0$ in [7]. They derived Hermitian self-orthogonal codes over F_4 as Gray images of linear and cyclic codes over $F_4 + uF_4$. In [9], X.Yin and W.Ma gave an existence condition of quantum codes which are derived from cyclic codes over finite ring $F_2 + uF_2 + u^2F_2$, $u^3 = 0$ with Lee metric. J.Qian gave a new method of constructing quantum error correcting codes from cyclic codes over finite ring $F_2 + vF_2$, $v^2 = v$, for arbitrary length n in [3].

This paper is organized as follows. In section 2, we give some basic knowledges about the finite ring R, cyclic code, dual code. In section 3, we define a new Gray map from R to F_2^3, Lee weights of elements of R and Lee distance in the linear codes over R. We show that if C is self orthogonal so is $\psi(C)$ and we obtain in the Gray images of cyclic codes over R. In section 4, a sufficient and necessary condition for cyclic code over R that contains its dual is given. The parameters of quantum error correcting codes are obtained from cyclic codes over R.

2 Preliminaries

Let R be the ring $F_2 + vF_2 + v^2F_2$ where $v^3 = v$ and $F_2 = \{0, 1\}$. R is a commutative ring with eight elements. R is semilocal ring with two maximal ideals and principal ideal ring with characteristic 2. It is not finite chain ring. It has six ideals. The ideal with 8 elements which generate the ring R is as follows

$$\langle 1 \rangle = \left\langle 1 + v + v^2 \right\rangle = R$$

The two maximal ideals with 4 elements are as follows

$$\begin{aligned} \langle v \rangle &= \left\langle v^2 \right\rangle = \left\{0, v, v^2, v + v^2\right\} \\ \langle 1 + v \rangle &= \left\{0, 1 + v, 1 + v^2, v + v^2\right\} \end{aligned}$$

The ideals with two elements are as follows

$$\begin{aligned} \left\langle 1 + v^2 \right\rangle &= \left\{0, 1 + v^2\right\} \\ \left\langle v + v^2 \right\rangle &= \left\{0, v + v^2\right\} \end{aligned}$$

The element 1 and $1 + v + v^2$ are two units of R. The zero divisors are $0, v, v^2, 1 + v, v + v^2, 1 + v + v^2$.

A linear code C over R length n is a $R-$submodule of R^n. An element of C is called a codeword.

For any $x = (x_0, x_1, ..., x_{n-1})$, $y = (y_0, y_1, ..., y_{n-1})$ the inner product is defined as

$$x.y = \sum_{i=0}^{n-1} x_i y_i$$

If $x.y = 0$ then x and y are said to be orthogonal. Let C be linear code of length n over R, the dual code of C

$$C^{\perp} = \{x : \forall y \in C, x.y = 0\}$$

which is also a linear code over R of length n. A code C is self orthogonal if $C \subseteq C^{\perp}$ and self dual if $C = C^{\perp}$.

A cyclic code C over R is a linear code with the property that if $c = (c_0, c_1, ..., c_{n-1}) \in C$ then $\sigma(C) = (c_{n-1}, c_0, ..., c_{n-2}) \in C$. A subset C of R^n is a linear cyclic code of length n iff it is polynomial representation is an ideal of $R[x] / \langle x^n - 1 \rangle$.

Let C be code over F_2 of length n and $\acute{c} = (\acute{c}_0, \acute{c}_1, ..., \acute{c}_{n-1})$ be a codeword of C. The Hamming weight of $\acute{c}$ is defined as $w_H(\acute{c}) = \sum_{i=0}^{n-1} w_H(\acute{c}_i)$ where $w_H(\acute{c}_i) = 1$ if $\acute{c}_i = 1$ and $w_H(\acute{c}_i) = 0$ if $\acute{c}_i = 0$. Hamming distance of C is defined as $d_H(C) = \min d_H(c, \acute{c})$, where for any $\acute{c} \in C$, $c \neq \acute{c}$ and $d_H(c, \acute{c})$ is Hamming distance between two codewords with $d_H(c, \acute{c}) = w_H(c - \acute{c})$.

Let $a \in F_2^{3n}$ with $a = (a_0, a_1, ..., a_{3n-1}) = \left(a^{(0)} \left| a^{(1)} \right| a^{(2)}\right)$, $a^{(i)} \in F_2^n$ for $i = 0, 1, 2$. Let φ be a map from F_2^{3n} to F_2^{3n} given by $\varphi(a) = \left(\sigma\left(a^{(0)}\right) \left|\sigma\left(a^{(1)}\right)\right| \sigma\left(a^{(2)}\right)\right)$ where σ is a cyclic shift from F_2^n to F_2^n given by $\sigma\left(a^{(i)}\right) = ((a^{(i,n-1)}), (a^{(i,0)}), ..., (a^{(i,n-2)}))$ for every $a^{(i)} = \left(a^{(i,0)}, ..., a^{(i,n-1)}\right)$ where $a^{(i,j)} \in F_2$, $j = 0, 1, ..., n-1$. A code of length $3n$ over F_2 is said to be quasi cyclic code of index 3 if $\varphi(C) = C$.

3 Gray Map And Gray Images Of Cyclic Codes Over R

Let $x = a + vb + v^2 c$ be an element of R where $a, b, c \in F_2$. We define Gray map ψ from R to F_2^3 by

$$\begin{array}{rcl} \psi & : & R \to F_2^3 \\ a + vb + v^2 c & \mapsto & \psi\left(a + vb + v^2 c\right) = (a, b, a + c) \end{array}$$

From definition, the Lee weight of elements of R are defined as follows

$$\begin{array}{rcll} w_L(0) & = & 0 & w_L(1 + v) = 3 \\ w_L(1) & = & 2 & w_L\left(1 + v^2\right) = 1 \\ w_L(v) & = & 1 & w_L\left(v + v^2\right) = 2 \\ w_L\left(v^2\right) & = & 1 & w_L\left(1 + v + v^2\right) = 2 \end{array}$$

Let C be a linear code over R of length n. For any codeword $c = (c_0, ..., c_{n-1})$ the Lee weight of c is defined as $w_L(c) = \sum_{i=0}^{n-1} w_L(c_i)$ and the Lee distance of C is defined as $d_L(C) = \min d_L(c, \acute{c})$, where for any $\acute{c} \in C$, $c \neq \acute{c}$ and $d_L(c, \acute{c})$ is Lee distance between two codewords with $d_L(c, \acute{c}) = w_L(c - \acute{c})$. Gray map ψ can be extended to map from R^n to F_2^{3n}.

Theorem 1 *The Gray map ψ is a weight preserving map from $(R^n, \text{Lee weight})$ to $\left(F_2^{3n}, \text{Hamming weight}\right)$. Moreover it is an isometry from R^n to F_2^{3n}.*

Proof. Similar to proof of theorem 3.5 in [9]. ■

Theorem 2 *If C is an (n, k, d_L) linear codes over R then $\psi(C)$ is a $(3n, k, d_H)$ linear codes over F_2, where $d_H = d_L$.*

Proof. Let $c = a_1 + vb_1 + v^2c_1$, $\acute{c} = a_2 + vb_2 + v^2c_2 \in R, \alpha \in F_2$ then

$$\begin{aligned}\psi(c + \acute{c}) &= \psi\left(a_1 + a_2 + v(b_1 + b_2) + v^2(c_1 + c_2)\right)\\ &= (a_1 + a_2, b_1 + b_2, a_1 + a_2 + c_1 + c_2)\\ &= (a_1, b_1, a_1 + c_1) + (a_2, b_2, a_2 + c_2)\\ &= \psi(c) + \psi(\acute{c})\end{aligned}$$

$$\begin{aligned}\psi(\alpha c) &= \psi\left(\alpha a_1 + v\alpha b_1 + v^2\alpha c_1\right)\\ &= (\alpha a_1, \alpha b_1, \alpha a_1 + \alpha c_1)\\ &= \alpha(a_1, b_1, a_1 + c_1)\\ &= \alpha\psi(c)\end{aligned}$$

so ψ is linear. As ψ is bijective then $|C| = |\psi(C)|$. From theorem 1 we have $d_H = d_L$. ■

Theorem 3 *If C is self orthogonal, so is $\psi(C)$.*

Proof. Let $c = a_1 + vb_1 + v^2c_1$, $\acute{c} = a_2 + vb_2 + v^2c_2$ where $a_1, b_1, c_1, a_2, b_2, c_2 \in F_2$.

$c.\acute{c} = a_1a_2 + v(a_1b_2 + b_1a_2 + b_1c_2 + c_1b_2) + v^2(a_1c_2 + b_1b_2 + c_1a_2 + c_1c_2)$, if C is self orthogonal,so we have $a_1a_2 = 0, a_1b_2 + b_1a_2 + b_1c_2 + c_1b_2 = 0, a_1c_2 + b_1b_2 + c_1a_2 + c_1c_2 = 0$. From

$$\begin{aligned}\psi(c).\psi(\acute{c}) &= (a_1, b_1, a_1 + c_1)(a_2, b_2, a_2 + c_2)\\ &= a_1a_2 + b_1b_2 + a_1a_2 + a_1c_2 + c_1a_2 + c_1c_2 = 0\end{aligned}$$

Therefore, we have $\psi(C)$ is self orthogonal. ■

Proposition 4 *Let ψ the Gray map from R^n to F_2^{3n} as follows*

$$\psi(r_0, r_1, ..., r_{n-1}) = (a_0, ..., a_{n-1}, b_0, ..., b_{n-1}, a_0 + c_0, ..., a_{n-1} + c_{n-1})$$

where $r_i = a_i + vb_i + v^2c_i$ for $i = 0, 1, ..., n-1$. Let σ be cyclic shift and let φ be a map from F_2^{3n} to F_2^{3n} as in the preliminaries. Then $\psi\sigma = \varphi\psi$.

Proof. Let $r_i = a_i + vb_i + v^2c_i$ be the elements of R for $i = 0, 1, ..., n-1$. We have $\sigma(r_0, r_1, ..., r_{n-1}) = (r_{n-1}, r_0, ..., r_{n-2})$. If we apply ψ, we have

$$\begin{aligned}\psi(\sigma(r_0, ..., r_{n-1})) &= \psi(r_{n-1}, r_0, ..., r_{n-2})\\ &= (a_{n-1}, ..., a_{n-2}, b_{n-1}, ..., b_{n-2}, a_{n-1} + c_{n-1}, ..., a_{n-2} + c_{n-2})\end{aligned}$$

On the other hand $\psi(r_0, ..., r_{n-1}) = (a_0, ..., a_{n-1}, b_0, ..., b_{n-1}, a_0 + c_0, ..., a_{n-1} + c_{n-1})$. If we apply φ, we have $\varphi(\psi(r_0, r_1, ..., r_{n-1})) = (a_{n-1}, ..., a_{n-2}, b_{n-1}, ..., b_{n-2}, a_{n-1} + c_{n-1}, ..., a_{n-2} + c_{n-2})$. Thus, $\psi\sigma = \varphi\psi$. ■

Proposition 5 *Let σ and φ be as above. A code C of length n over R is cyclic code if and only if $\psi(C)$ is quasi cyclic code of index 3 and length $3n$ over F_2.*

Proof. Suppose C is cyclic code. Then $\sigma(C) = C$. If we apply ψ, we have $\psi(\sigma(C)) = \psi(C)$. From proposition 4, $\psi(\sigma(C)) = \varphi(\psi(C)) = \psi(C)$. Hence, $\psi(C)$ is a quasi cyclic code of index 3. Conversely, if $\psi(C)$ is a quasi cyclic code of index 3, then $\varphi(\psi(C)) = \psi(C)$. From proposition 4, we have $\varphi(\psi(C)) = \psi(\sigma(C)) = \psi(C)$. Since ψ is injective, it follows that $\sigma(C) = C$. ■

We know that any linear code over R of length n is permutation equivalent to a code with generator matrix of the form

$$G = \begin{bmatrix} I_{k_1} & A_1 & A_2 & A_{31} + vA_{32} & A_{41} + vA_{42} & A_5 \\ 0 & vI_{k_2} & B_1 & B_{21} + vB_{22} & vB_{31} & B_4 \\ 0 & 0 & (1+v) I_{k_3} & (1+v) C_1 & (1+v) C_2 & C_3 \\ 0 & 0 & 0 & \left(v + v^2\right) I_{k_4} & \left(v + v^2\right) D_1 & D \\ 0 & 0 & 0 & 0 & \left(1 + v^2\right) I_{k_5} & E \end{bmatrix}$$

where $I_{k_1}, I_{k_2}, I_{k_3}, I_{k_4}, I_{k_5}$ are all unit matrices with order k_1, k_2, k_3, k_4, k_5 respectively. Let $k = k_1 + k_2 + k_3 + k_4 + k_5$ [8]. $A_5 = A_{51} + vA_{52} + v^2A_{53}$, $B_4 = vB_{41} + v^2B_{42}$, $C_1 = C_{11} + vC_{12} + v^2C_{13}$, $C_3 = (1+v)\left(C_{31} + vC_{32} + v^2C_{33}\right)$, $D = \left(v + v^2\right) D_2$, $E = \left(1 + v^2\right) E'$ where $A_1, A_2, A_{31}, A_{32}, A_{41}, A_{42}, A_{51}, A_{52}, A_{53}, B_1, B_{21}, B_{22}, B_{31}, B_{41}, B_{42}, C_{11}, C_{12}, C_{13}, C_2, C_{31}, C_{32}, C_{33}, D_1, D_2, E'$ are matrices over F_2 in [8].

Let C be a linear code of length n over R. Define

$$\begin{aligned} C_1 &= \left\{a \in F_2^n : \exists b, c \in F_2^n, a + vb + v^2c \in C\right\} \\ C_2 &= \left\{b \in F_2^n : \exists a, c \in F_2^n, a + vb + v^2c \in C\right\} \\ C_3 &= \left\{a + c \in F_2^n : \exists b \in F_2^n, a + vb + v^2c \in C\right\} \end{aligned}$$

Obviously, C_1, C_2 and C_3 are binary linear codes.

We denote that $A \otimes B \otimes C = \{(a, b, c) : a \in A, b \in B, c \in C\}$ and $A \oplus B \oplus C = \{a + b + c : a \in A, b \in B, c \in C\}$.

Theorem 6 *Let C be a linear code of length n over R. Then $\psi(C) = C_1 \otimes C_2 \otimes C_3$ and $|C| = |C_1||C_2||C_3|$.*

Proof. For any $(a_0, a_1, ..., a_{n-1}, b_0, b_1, ..., b_{n-1}, a_0 + c_0, a_1 + c_1, ..., a_{n-1} + c_{n-1}) \in \psi(C)$. Let $d_i = v(a_i + b_i + c_i) + (1+v)(a_i + b_i) + \left(1 + v^2\right) c_i$, $i = 0, 1, ..., n-1$. Since ψ is a bijection $d = (d_0, d_1, ..., d_{n-1}) \in C$. By definitions of C_1, C_2 and C_3 we have $(a_0, a_1, ..., a_{n-1}) \in C_1$, $(b_0, b_1, ..., b_{n-1}) \in C_2$, $(a_0 + c_0, a_1 + c_1, ..., a_{n-1} + c_{n-1}) \in C_3$. So, $(a_0, a_1, ..., a_{n-1}, b_0, b_1, ..., b_{n-1}, a_0 + c_0, a_1 + c_1, ..., a_{n-1} + c_{n-1}) \in$

$C_1 \otimes C_2 \otimes C_3$. That is $\psi(C) \subseteq C_1 \otimes C_2 \otimes C_3$. On the other hand, for any $(a_0, a_1, ..., a_{n-1}, \quad b_0, b_1, ..., b_{n-1}, a_0+c_0, a_1+c_1, ..., a_{n-1}+c_{n-1}) \in C_1 \otimes C_2 \otimes C_3$ where $(a_0, a_1, ..., a_{n-1}) \in C_1$, $(b_0, b_1, ..., b_{n-1}) \in C_2$, $(a_0 + c_0, a_1 + c_1, ..., a_{n-1} + c_{n-1}) \in C_3$. There are $x = (a_0, a_1, ..., a_{n-1})$, $y = (b_0, b_1, ..., b_{n-1})$, $z = (c_0, c_1, ..., c_{n-1}) \in C$ such that $x_i = a_i + vp_i$, $y_i = b_i + (1+v) q_i$, $z_i = (a_i + c_i) + \left(1 + v^2\right) r_i$ where $p_i, q_i, r_i \in F_2$ and $0 \leq i \leq n-1$. Since C is linear we have $d = vx + (1+v) y + \left(1+v^2\right) z = v(a+b+c) + (1+v)(a+c) + \left(1+v^2\right) c \in C$. It follows then $\psi(d) = (a_0, a_1, ..., a_{n-1}, b_0, b_1, ..., b_{n-1}, a_0 + c_0, a_1 + c_1, ..., a_{n-1} + c_{n-1})$, which gives $C_1 \otimes C_2 \otimes C_3 \subseteq \psi(C)$. Therefore, $\psi(C) = C_1 \otimes C_2 \otimes C_3$. The second result is easy to verify. ■

Corollary 7 *If $\psi(C) = C_1 \otimes C_2 \otimes C_3$, then $C = vC_1 \oplus (1+v) C_2 \oplus \left(1+v^2\right) C_3$.*

Moreover,

$$G_1 = \begin{bmatrix} I_{k_1} & A_1 & A_3 & A_{31} & A_{41} & A_{51} \\ 0 & 0 & B_1 & B_{21} & 0 & 0 \\ 0 & 0 & I_{k_3} & C_1 & C_2 & C_{31} \\ 0 & 0 & 0 & 0 & I_{k_5} & E' \end{bmatrix}$$

$$G_2 = \begin{bmatrix} 0 & 0 & 0 & A_{32} & A_{42} & A_{52} \\ 0 & I_{k_2} & 0 & B_{22} & B_{31} & B_{41} \\ 0 & 0 & I_{k_3} & C_{11}+C_{12}+C_{13} & C_2 & C_{31}+C_{32}+C_{33} \\ 0 & 0 & 0 & I_{k_4} & D_1 & D_2 \end{bmatrix}$$

$$G_3 = \begin{bmatrix} I_{k_1} & A_1 & A_2 & A_{31} & A_{41} & A_{51}+A_{53} \\ 0 & 0 & B_1 & B_{21} & 0 & B_{43} \\ 0 & 0 & I_{k_3} & C_{11}+C_{12}+C_{13} & C_2 & C_{31}+C_{32}+C_{33} \\ 0 & 0 & 0 & I_{k_4} & D_1 & D_2 \end{bmatrix}$$

It is easy to see that $|C| = |C_1||C_2||C_3| = 2^{3(k_1+k_2+k_3)+2k_4+k_5} = 8^{k_1+k_2+k_3} 4^{k_4} 2^{k_5}$.

Corollary 8 *If G_1, G_2,and G_3 are generator matrices of binary linear codes C_1, C_2 and C_3 respectively, then the generator matrix of C is*

$$G = \begin{bmatrix} vG_1 \\ (1+v) G_2 \\ \left(1+v^2\right) G_3 \end{bmatrix}$$

■

Let d_L minimum Lee weight of linear code C over R. Then $d_L = \min\{d_H(C_1), d_H(C_2), d_H(C_3)\}$ where $d_H(C_i)$ denotes the minimum Hamming weights of binary codes C_1, C_2 and C_3.

4 Quantum Codes From Cyclic Codes Over R

Theorem 9 (*CSS Construction*) *Let C and $\acute{C}$ be two binary codes with parameters $[n, k_1, d_1]$ and $[n, k_2, d_2]$, respectively. If $C^{\perp} \subseteq \acute{C}$, then an $[[n, k_1 + k_2 -$*

$n, \min\{d_1, d_2\}]]$ *quantum code can be constructed. Especially, if* $C^{\perp} \subseteq C$*, then there exists an* $[[n, 2k_1 - n, d_1]]$ *quantum code.*

Proposition 10 *Let* $C = vC_1 \oplus (1+v) C_2 \oplus \left(1+v^2\right) C_3$ *be a linear code over R.Then C is a cyclic code over R iff* C_1, C_2 *and* C_3 *are binary cyclic codes.*

Proposition 11 *Suppose* $C = vC_1 \oplus (1+v) C_2 \oplus \left(1+v^2\right) C_3$ *is a cyclic code of length n over R. Then* $C = \left\langle vf_1, (1+v) f_2, \left(1+v^2\right) f_3 \right\rangle$ *and* $|C| = 2^{3n-(\deg f_1 + \deg f_2 + \deg f_3)}$ *where* f_1, f_2 *and* f_3 *generator polynomials of* C_1, C_2 *and* C_3 *respectively.*

Proposition 12 *Suppose C is a cyclic code of length n over R, then there is a unique polynomial* $f(x)$ *such that* $C = \langle f(x) \rangle$ *and* $f(x) \mid x^n - 1$ *where* $f(x) = vf_1(x) + (1+v) f_2(x) + \left(1+v^2\right) f_3(x)$.

Proposition 13 *If* $C = vC_1 \oplus (1+v) C_2 \oplus \left(1+v^2\right) C_3$ *is a cyclic code of length n over R, then* $C^{\perp} = \left\langle vh_1^* + (1+v) h_2^* + \left(1+v^2\right) h_3^* \right\rangle$ *and* $\left|C^{\perp}\right| = 2^{\deg f_1 + \deg f_2 + \deg f_3}$ *where for* $i = 1, 2, 3$*,* h_i^* *are the reciprocal polynomials of* h_i *i.e.,* $h_i(x) = (x^n - 1)/f_i(x)$, $h_i^*(x) = x^{\deg h_i} h_i\left(x^{-1}\right)$ *for* $i = 1, 2, 3$.

Lemma 14 *A binary linear cyclic code C with generator polynomial* $f(x)$ *contains its dual code iff*

$$x^n - 1 \equiv 0 \pmod{ff^*}$$

where f^* is the reciprocal polynomial of f.

Theorem 15 *Let* $C = \left\langle vf_1 + (1+v) f_2 + \left(1+v^2\right) f_3 \right\rangle$ *be a cyclic code of length n over R. Then* $C^{\perp} \subseteq C$ *iff* $x^n - 1 \equiv 0 \pmod{f_i f_i^*}$ *for* $i = 1, 2, 3$.

Proof. Let $x^n - 1 \equiv 0 \pmod{f_i f_i^*}$ for $i = 1, 2, 3$. Then $C_1^{\perp} \subseteq C_1, C_2^{\perp} \subseteq C_2, C_3^{\perp} \subseteq C_3$. By using $vC_1^{\perp} \subseteq vC_1$, $(1+v) C_2^{\perp} \subseteq (1+v) C_2$, $\left(1+v^2\right) C_3^{\perp} \subseteq \left(1+v^2\right) C_3$. We have $vC_1^{\perp} \oplus (1+v) C_2^{\perp} \oplus \left(1+v^2\right) C_3^{\perp} \subseteq vC_1 \oplus (1+v) C_2 \oplus \left(1+v^2\right) C_3$. So, $\left\langle vh_1^* + (1+v) h_2^* + \left(1+v^2\right) h_3^* \right\rangle \subseteq \left\langle vf_1, (1+v) f_2, \left(1+v^2\right) f_3 \right\rangle$. That is $C^{\perp} \subseteq C$.

Conversely, if $C^{\perp} \subseteq C$, then $vC_1^{\perp} \oplus (1+v) C_2^{\perp} \oplus \left(1+v^2\right) C_3^{\perp} \subseteq vC_1 \oplus (1+v) C_2 \oplus \left(1+v^2\right) C_3$. By thinking $\bmod v, \bmod (1+v), \bmod \left(1+v^2\right)$ respectively we have $C_i^{\perp} \subseteq C_i$ for $i = 1, 2, 3$. Therefore, $x^n - 1 \equiv 0 \pmod{f_i f_i^*}$ for $i = 1, 2, 3$. ■

Corollary 16 $C = vC_1 \oplus (1+v) C_2 \oplus \left(1+v^2\right) C_3$ *is a cyclic code of length n over R. Then* $C^{\perp} \subseteq C$ *iff* $C_i^{\perp} \subseteq C_i$ *for* $i = 1, 2, 3$.

Example 17 *Let* $n = 21, R = F_2 + vF_2 + v^2 F_2$

$$\begin{aligned} x^{21} - 1 &= (x+1)(x^2+x+1)(x^3+x^2+1)(x^3+x+1)(x^6+x^4+x^2+x+1) \\ &\quad (x^6+x^5+x^4+x^2+1) \end{aligned}$$

in $F_2[x]$.

Let $f_1 = x+1$, $f_2 = x^2+x+1$, $f_3 = x^3+x^2+1$, $f_4 = x^3+x+1$, $f_5 = x^6+x^4+x^2+x+1$, $f_6 = x^6+x^5+x^4+x^2+1$. Hence,

$f_1^* = x+1 = f_1$, $f_2^* = x^2+x+1 = f_2$, $f_3^* = x^3+x+1 = f_4$, $f_4^* = x^3+x^2+1 = f_3$, $f_5^* = x^6+x^5+x^4+x^2+1 = f_6$, $f_6^* = x^6+x^4+x^2+x+1 = f_5$.

Let $C = \langle vf_i, (1+v)f_i, (1+v^2)f_i \rangle$ for $i = 3,4,5,6$. Obviously $x^n - 1$ is divisibly by $f_i f_i^*$, for i. Thus we have $C^\perp \subseteq C$.

Using Theorem 9 and Theorem 15 we can construct quantum codes.

Theorem 18 *Let $C = vC_1 \oplus (1+v)C_2 \oplus (1+v^2)C_3$ be a cyclic code of arbitrary length n over R with type $8^{k_1+k_2+k_3}4^{k_4}2^{k_5}$. If $C_i^\perp \subseteq C_i$ where $i = 1,2,3$ then $C^\perp \subseteq C$ and there exists a quantum error-correcting code with parameters $[[3n, 6(k_1+k_2+k_3)+4k_4+2k_5-3n, d_L]]$ where d_L is the minimum Lee weights of C.*

5 Examples

Example 19 *Let $n = 8$*

$$x^8 - 1 = (x+1)(x+1)(x+1)(x+1)(x+1)(x+1)(x+1)(x+1)$$

in $F_2[x]$. Let $f_1(x) = f_2(x) = f_3(x) = x^3+x^2+x+1$.

Thus $C = \langle vf_1, (1+v)f_2, (1+v^2)f_3 \rangle = \langle v^2(x^3+x^2+x+1) \rangle$. C is a linear code of length 8 and minimum Lee weight $d_L = 2$. The dual code $C^\perp = \langle vh_1^*, (1+v)h_2^*, (1+v^2)h_3^* \rangle = \langle v^2(x^5+x^4+x+1) \rangle$ can be obtained. Clearly, $C^\perp \subseteq C$. Hence we obtain a quantum code with parameters $[[24,6,2]]$.

Example 20 *Let $n = 8$*

$$x^8 - 1 = (x+1)(x+1)(x+1)(x+1)(x+1)(x+1)(x+1)(x+1)$$

in $F_2[x]$. Let $f_1(x) = f_2(x) = f_3(x) = x^2+1$.

Thus $C = \langle vf_1, (1+v)f_2, (1+v^2)f_3 \rangle = \langle v^2(x^2+1) \rangle$. C is a linear code of length 8 and minimum Lee weight $d_L = 2$.

The dual code $C^\perp = \langle vh_1^*, (1+v)h_2^*, (1+v^2)h_3^* \rangle = \langle v^2(x^6+x^4+x^2+1) \rangle$ can be obtained. Clearly, $C^\perp \subseteq C$. Hence we obtain a quantum code with parameters $[[24,12,2]]$. Let $f_1(x) = f_2(x) = f_3(x) = x+1$.Thus we obtain a quantum code with parameters $[[24,18,2]]$.

Example 21 *Let $n = 7$*

$$x^7 - 1 = (x+1)\left(x^3+x+1\right)\left(x^3+x^2+1\right)$$

in $F_2[x]$. Let $f_1(x) = f_2(x) = f_3(x) = x^3 + x + 1$.

Thus $C = \left\langle vf_1, (1+v) f_2, \left(1+v^2\right) f_3 \right\rangle = \left\langle v^2\left(x^3+x+1\right)\right\rangle$. C is a linear code of length 7 and minimum Lee weight $d_L = 3$.The dual code $C^\perp = \left\langle vh_1^*, (1+v) h_2^*, \left(1+v^2\right) h_3^* \right\rangle = \left\langle v^2\left(x^4+x^2+x+1\right)\right\rangle$ can be obtained. Clearly, $C^\perp \subseteq C$. Hence we obtain a quantum code with parameters $[[21, 3, 3]]$.

Example 22 *Let* $n = 15$

$$x^{15}-1 = (x+1)\left(x^2+x+1\right)\left(x^4+x+1\right)\left(x^4+x^3+1\right)\left(x^4+x^3+x^2+x+1\right)$$

in $F_2[x]$. Let $f_1(x) = f_2(x) = f_3(x) = x^4 + x + 1$.

Thus $C = \left\langle vf_1, (1+v) f_2, \left(1+v^2\right) f_3 \right\rangle = \left\langle v^2\left(x^4+x+1\right)\right\rangle$. C is a linear code of length 15 and minimum Lee weight $d_L = 3$.The dual code $C^\perp = \left\langle vh_1^*, (1+v) h_2^*, \left(1+v^2\right) h_3^* \right\rangle$ can be obtained. Clearly, $C^\perp \subseteq C$. Hence we obtain a quantum code with parameters $[[45, 21, 3]]$.

Example 23 *Let* $n = 16$

$$x^{16} - 1 = (x+1)^{16}$$

in $F_2[x]$. Let $f_1(x) = f_2(x) = f_3(x) = x^3 + x^2 + x + 1$.

Thus $C = \left\langle vf_1, (1+v) f_2, \left(1+v^2\right) f_3 \right\rangle = \left\langle v^2\left(x^3+x^2+x+1\right)\right\rangle$. C is a linear code of length 16 and minimum Lee weight $d_L = 2$.The dual code $C^\perp = \left\langle vh_1^*, (1+v) h_2^*, \left(1+v^2\right) h_3^* \right\rangle = \left\langle v^2\left(x^{13}+x^{12}+x^9+x^8+x^5+x^4+x+1\right)\right\rangle$ can be obtained. Clearly, $C^\perp \subseteq C$. Hence we obtain a quantum code with parameters $[[48, 30, 2]]$. Let $f_1(x) = f_2(x) = f_3(x) = x^4 + 1$. Thus we obtain a quantum code with parameters $[[48, 24, 2]]$.

Example 24 *Let* $n = 21$,

$$\begin{aligned} x^{21} - 1 \;\; = \;\; & (x+1)(x^2+x+1)(x^3+x^2+1)(x^3+x+1)(x^6+x^4+x^2+x+1) \\ & (x^6+x^5+x^4+x^2+1) \end{aligned}$$

in $F_2[x]$. Let $f_1(x) = f_2(x) = f_3(x) = x^6 + x^5 + x^4 + x^2 + 1$.

Thus $C = \left\langle vf_1, (1+v) f_2, \left(1+v^2\right) f_3 \right\rangle = \left\langle v^2\left(x^4+x+1\right)\right\rangle$. C is a linear code of length 21 and minimum Lee weight $d_L = 3$.The dual code $C^\perp = \left\langle vh_1^*, (1+v) h_2^*, \left(1+v^2\right) h_3^* \right\rangle$ can be obtained. Clearly, $C^\perp \subseteq C$. Hence we obtain a quantum code with parameters $[[63, 27, 3]]$. Let $f_1(x) = f_2(x) = f_3(x) = x^3 + x^2 + 1$. Thus we obtain a quantum code with parameters $[[63, 45, 3]]$.

Acknowledgement

We would like to thank S.T.Dougherty for his many suggestions which helped us.

6 References

[1] A.M.Steane,Simple quantum error correcting codes, Phys. Rev. A, $54\,(1996)$, $4741-4751$.

[2] A.R.Calderbank, E.M.Rains, P.M.Shor, N.J.A.Sloane, Quantum error correction via codes over $GF\,(4)$, IEEE Trans. Inf. Theory, $44\,(1998)$, $1369-1387$.

[3] J.Qian, Quantum codes from cyclic codes over $F_2 + vF_2$, Journal of Inform.& computational Science $10:6\,(2013)$, $1715-1722$.

[4] J.Qian, W.Ma, W.Gou, Quantum codes from cyclic codes over finite ring, Int. J. Quantum Inform., $7\,(2009)$, $1277-1283$.

[5] P.W.Shor,Scheme for reducing decoherence in quantum memory, Phys. Rev. A, $52\,(1995)$, $2493-2496$.

[6] S.Zhu, Y.Wang, M.Shi, Some Results On Cyclic Codes Over $F_2 + vF_2$, IEEE Transactions on Information Theory,vol.56, No.4, 2010

[7] X.Kai,S.Zhu, Quaternary construction bof quantum codes from cyclic codes over $F_4 + uF_4$, Int. J. Quantum Inform., $9\,(2011)$, $689-700$.

[8] X.Xu, S.Yun, Y.Xiong, Generator Matrix of The Linear Codes And Gray Images Over Ring $F_2 + vF_2 + v^2F_2$, Journal of Applied Sciences $13\,(4)\,(2013):650-653$.

[9] X.Yin, W.Ma, Gray Map And Quantum Codes Over The Ring $F_2+uF_2+u^2F_2$, International Joint Conferences of IEEE TrustCom-11,2011.